

A TWO PARAMETER DEFORMATION OF THE $SU(1/1)$ SUPERALGEBRA AND THE XY QUANTUM CHAIN IN A MAGNETIC FIELD

Haye Hinrichsen

*Universität Bonn, Physikalisches Institut
Nussallee 12, W-5300 Bonn 1, FRG*

Vladimir Rittenberg*

*Theory Division, CERN
CH-1211 Geneva 23, Switzerland*

Abstract

We show that the XY quantum chain in a magnetic field is invariant under a two parameter deformation of the $SU(1/1)$ superalgebra. One is led to an extension of the braid group and the Hecke algebra which reduce to the known ones when the two parameter coincide. The physical significance of the two parameters is discussed.

CERN-TH.6299/91
October 1991

*Permanent address: Universität Bonn, Physikalisches Institut, Nussallee 12, W-5300 Bonn 1, FRG

There were several attempts to extend the one-parameter quantum algebras to multiparameter ones [1]. As shown however by Reshetikhin [2] the link polynomials depend only on one parameter. One can state this result in a different way: if one has a one-dimensional quantum chain which is invariant under a multiparameter quantum algebra, one can do a similarity transformation which eliminates all the parameters but one. As will be shown in this paper, the situation is different in the case of quantum superalgebras. We will start with a physical example. Consider the quantum chain

$$H = \Omega_q \sum_{i=1}^L \sigma_i^z + \frac{\Omega_\eta}{2} \sum_{i=1}^{L-1} [(1+u)\sigma_i^x \sigma_{i+1}^x + (1-u)\sigma_i^y \sigma_{i+1}^y] + B + S, \quad (1)$$

where σ^x , σ^y and σ^z are Pauli matrices inserted in the i -th position of the Kronecker product

$$\begin{aligned} \sigma_i^k &= \mathbf{1} \otimes \mathbf{1} \otimes \dots \otimes \underbrace{\sigma_i^k}_{\mathbf{i}} \otimes \dots \otimes \mathbf{1} \otimes \mathbf{1} & (i = 1, 2, \dots, L) \\ [\sigma_i^k, \sigma_j^l] &= 0. & (i \neq j) \end{aligned} \quad (2)$$

Ω_q , Ω_η and u are parameters, B and S are boundary and surface terms respectively. This chain appears in the domain wall theory of two-dimensional commensurate-incommensurate phase transitions [3] and in Glauber's kinetic Ising model [4]. We will first make an important change of notations, choose $B = 0$ (no periodic boundary conditions!) and fix S by

$$\begin{aligned} \Omega_q &= \frac{q + q^{-1}}{2}, & \Omega_\eta &= \frac{\eta + \eta^{-1}}{2}, & u &= \frac{\eta - \eta^{-1}}{\eta + \eta^{-1}} \\ S &= \frac{1}{2} (-q^{-1} \sigma_1^z - q \sigma_L^z). \end{aligned} \quad (3)$$

With this change of notations we have

$$\begin{aligned} H &= H(q, \eta) = \sum_{i=1}^{L-1} H_i(q, \eta) \\ H_i(q, \eta) &= \frac{1}{2} [\eta \sigma_i^x \sigma_{i+1}^x + \eta^{-1} \sigma_i^y \sigma_{i+1}^y + q \sigma_i^z + q^{-1} \sigma_{i+1}^z]. \end{aligned} \quad (4)$$

A detailed discussion of the properties of the chain given by eq. (4) will be given elsewhere [5], here we are going to mention only a few. First, there are the symmetry properties

$$H(q, \eta) \doteq H(q^{-1}, \eta) \doteq H(q, \eta^{-1}) \doteq H(\eta, q). \quad (5)$$

The "equality" among the Hamiltonians implies that the spectra are identical. The first two equalities are obvious but not the last one which reminds of duality transformations of quantum chains [6]. In the continuum limit, one has the following phase structure [3, 5]:

$\Omega_q \leq 1, \quad \Omega_\eta \leq 1:$	massless-incommensurate
$\Omega_q \leq 1, \quad \Omega_\eta > 1 \text{ or } \Omega_q > 1, \quad \Omega_\eta \leq 1:$	massive incommensurate
$\Omega_q > 1, \quad \Omega_\eta > 1, \quad \Omega_q \neq \Omega_\eta:$	massive
$\Omega_q = \Omega_\eta, \quad (\Omega_q > 1) :$	critical Ising type
$\Omega_q = \Omega_\eta = 1 :$	Pokrovsky-Talapov phase transition

It is by now clear that the properties of the chain depend on both parameters q and η . We now perform a Jordan-Wigner transformation. First write $\sigma_j^z = -i\sigma_j^x\sigma_j^y$ and next define

$$\begin{aligned} \tau_j^{x,y} &= \exp\left(\frac{i\pi}{2} \sum_{k=1}^{j-1} (\sigma_k^z + 1)\right) \sigma_j^{x,y} \\ \{\tau_i^x, \tau_j^x\} &= \{\tau_i^y, \tau_j^y\} = \{\tau_i^x, \tau_j^y\} = 0. \end{aligned} \quad (i \neq j) \quad (6)$$

Using Eq. (4) and (6) we get

$$H_j = \frac{1}{2}[\eta\tau_j^x\tau_{j+1}^x + \eta^{-1}\tau_j^y\tau_{j+1}^y - iq\tau_j^x\tau_j^y - iq^{-1}\tau_{j+1}^x\tau_{j+1}^y]. \quad (7)$$

We now observe the following important identity

$$[T^X, H(q, \eta)] = [T^Y, H(q, \eta)] = 0 \quad (8)$$

with

$$\begin{aligned} T^X &= \Delta(\tau^x) = \sum_{j=1}^L \alpha^{\frac{1-L}{2}} \sum_{j=1}^L \alpha^{j-1} \tau_j^x \\ T^Y &= \Delta(\tau^y) = \sum_{j=1}^L \beta^{\frac{1-L}{2}} \sum_{j=1}^L \beta^{j-1} \tau_j^y \end{aligned} \quad (9)$$

$$\{T^X, T^X\} = 2[L]_\alpha, \quad \{T^Y, T^Y\} = 2[L]_\beta, \quad \{T^X, T^Y\} = 0, \quad (10)$$

where L is the length of the chain and

$$\alpha = -\frac{q}{\eta}, \quad \beta = -q\eta, \quad [L]_\lambda = \frac{\lambda^L - \lambda^{-L}}{\lambda - \lambda^{-1}}. \quad (11)$$

The equalities (8) come from the existence of a fermionic zero mode for a q and η . The equations (10) together with the coproduct (9) give a representation of a Hopf algebra. Before we prove this statement let us consider the case $\alpha = \beta = -q$. We first notice that in this case $S^z = \frac{1}{2} \sum_{i=1}^L \sigma_i^z$ also commutes with $H(q, \eta)$. We now remind the reader the $U_\alpha[SU(1/1)]$ algebra [7]. With $A^\pm = \frac{1}{2}(T^X \pm iT^Y)$ we have

$$\begin{aligned} \{A^\pm, A^\pm\} &= 0, & \{A^+, A^-\} &= [E]_\alpha, & [S^z, A^\pm] &= \pm A^\pm \\ [E, S^z] &= [E, A^\pm] = 0 \end{aligned} \quad (12)$$

with the coproduct

$$\begin{aligned} \Delta(\alpha, A^\pm) &= \alpha^{E/2} \otimes A^\pm + A^\pm \otimes \alpha^{-E/2} \\ \Delta(\alpha, S^z) &= S^z \otimes \mathbf{1} + \mathbf{1} \otimes S^z \\ \Delta(\alpha, E) &= E \otimes \mathbf{1} + \mathbf{1} \otimes E. \end{aligned} \quad (13)$$

The fermionic representations correspond to take $E = \mathbf{1}$, $S^z = \frac{1}{2}\sigma^z$, $A^\pm = a^\pm$ and $\{a^+, a^-\} = 1$ in Eq (13). In this representation E in Eq. (12) is equal to L (the number of sites). Comparing now (12), (13) with Eqs. (9,10) we observe [8] that the quantum chain (4) with $\eta = 1$ is invariant under $U_\alpha[SU(1/1)]$ transformations. It was also shown by Saleur [8] that the quantities $U_j = \Delta_q - H_j(q, 1)$ are the generators of the Heck algebra

$$\begin{aligned} U_j^2 &= 2 \Delta_q U_j \\ U_j U_{j\pm 1} U_j - U_j &= U_{j\pm 1} U_j U_{j\pm 1} - U_{j\pm 1} \\ U_i U_{i\pm j} &= U_{i\pm j} U_i. \end{aligned} \quad (j \neq 1) \quad (14)$$

Actually they correspond to a quotient of this algebra since the generators satisfy also the relations [9]

$$U_j U_{j+2} U_{j+1} (2\Omega_q - U_j) (2\Omega_q - U_{j+2}) = 0. \quad (15)$$

The generators $\check{R}_j = \frac{q-q^{-1}}{2} + H_j(q, 1)$ satisfy the braiding relations

$$\check{R}_j \check{R}_{j\pm 1} \check{R}_j = \check{R}_{j\pm 1} \check{R}_j \check{R}_{j\pm 1} \quad (16)$$

with

$$\check{R}_j^2 = (q - q^{-1}) \check{R}_j + 1. \quad (17)$$

Considering the matrices $R_j = P \check{R}_j$ (P is the graded permutation operator) we have (see Eq. (13))

$$R \Delta(\alpha) R^{-1} = \Delta(\alpha^{-1}). \quad (18)$$

We now consider the case $\eta \neq 1$. As suggested by Eqs. (9,10) we define the two parameter deformation of the $SU(1/1)$ algebra as follows:

$$\begin{aligned} \{T^X, T^X\} &= 2 [E]_\alpha, & \{T^Y, T^Y\} &= 2 [E]_\beta \\ \{T^X, T^Y\} &= 0 & [E, T^X] &= [E, T^Y] = 0 \end{aligned} \quad (19)$$

with the coproduct

$$\begin{aligned}
\Delta(\alpha, \beta; T^X) &= \alpha^{E/2} \otimes T^X + T^X \otimes \alpha^{-E/2} \\
\Delta(\alpha, \beta; T^Y) &= \beta^{E/2} \otimes T^Y + T^Y \otimes \beta^{-E/2} \\
\Delta(\alpha, \beta; E) &= E \otimes \mathbf{1} + \mathbf{1} \otimes E.
\end{aligned} \tag{20}$$

Notice that S^z does not appear in the algebra anymore. We denote this quantum algebra by $U_{\alpha, \beta}[SU(1/1)]$. It is a Hopf algebra for the same reasons as the $U_\alpha[SU(1/1)]$. If we take the fermionic representations $E = 1$, $\tau^x = (a^+ + a^-)$, $\tau^y = -i(a^+ - a^-)$, from Eq. (20) we derive Eqs. (9,10). The quantum chain $H(q, \eta)$ is thus invariant under the quantum algebra $U_{\alpha, \beta}[SU(1/1)]$. We would like to see what replaces the relations (14-18) when we have two parameters. We first notice a remarkable identity satisfied by the $H_j(q, \eta)$

$$[H_j H_{j\pm 1} H_j - H_{j\pm 1} H_j H_{j\pm 1} + (\nu - 1)(H_j - H_{j\pm 1})] (H_j - H_{j\pm 1}) = \mu \tag{21}$$

$$H_j^2 = \nu,$$

where

$$\begin{aligned}
\nu &= \left(\frac{\alpha + \alpha^{-1}}{2} \right) \left(\frac{\beta + \beta^{-1}}{2} \right) = \left(\frac{q + q^{-1}}{2} \right)^2 + \left(\frac{\eta + \eta^{-1}}{2} \right)^2 - 1 \\
\mu &= \left(\frac{\alpha + \alpha^{-1}}{2} - \frac{\beta + \beta^{-1}}{2} \right)^2 = 4 \left(\frac{q - q^{-1}}{2} \right)^2 \left(\frac{\eta - \eta^{-1}}{2} \right)^2.
\end{aligned} \tag{22}$$

We can now define a generalised Hecke algebra taking

$$\begin{aligned}
U_i &= \sqrt{\nu} - H_i(p, q) \\
(U_i U_{i\pm 1} U_i - U_{i\pm 1} U_i U_{i\pm 1} - U_i + U_{i\pm 1}) (U_i - U_{i\pm 1}) &= \mu \\
U_i^2 &= 2\sqrt{\nu} U_i.
\end{aligned} \tag{23}$$

Notice that when $\eta = 1$, $\mu = 0$ and we are back to the original Hecke algebra. We did not have the patience to find the equivalent of Eq. (15) which gives the quotient of the generalised Hecke algebra (23) corresponding to the chain given by Eq. (4). Another quotient is however suggested by the structure of Eq. (23):

$$(U_i U_{i\pm 1} U_i - U_i) (U_i - U_{i\pm 1}) = \frac{\mu}{2}. \tag{24}$$

For $\mu = 0$ one gets in this case the Temperley-Lieb algebra. We now turn our attention to the generalised braid group algebra. We take $\check{R}_i = H_i(q, \eta) + \sqrt{\nu - 1}$ and get

$$(\check{R}_i \check{R}_{i\pm 1} \check{R}_i - \check{R}_{i\pm 1} \check{R}_i \check{R}_{i\pm 1}) (\check{R}_i - \check{R}_{i\pm 1}) = \mu \tag{25}$$

with

$$\check{R}_i^2 = 1 + \sqrt{\nu - 1} \check{R}_i. \quad (26)$$

In the basis where the σ_i^z are diagonal (see Eq. (4)) we have

$$\check{R}_i = \begin{pmatrix} \sqrt{\nu - 1} + \frac{q+q^{-1}}{2} & 0 & 0 & \frac{\eta-\eta^{-1}}{2} \\ 0 & \sqrt{\nu - 1} - \frac{q-q^{-1}}{2} & \frac{\eta+\eta^{-1}}{2} & 0 \\ 0 & \frac{\eta+\eta^{-1}}{2} & \sqrt{\nu - 1} + \frac{q+q^{-1}}{2} & 0 \\ \frac{\eta-\eta^{-1}}{2} & 0 & 0 & \sqrt{\nu - 1} - \frac{q+q^{-1}}{2} \end{pmatrix}. \quad (27)$$

We take the graded permutation matrix P

$$P = \begin{pmatrix} 1 & & & \\ & 0 & 1 & \\ & 1 & 0 & \\ & & & -1 \end{pmatrix} \quad (28)$$

and define the matrix $R_i = P \check{R}_i$. We now write the coproduct (20) in the original language of Pauli matrices

$$\begin{aligned} \Delta(\alpha, \beta; T^X) &= \alpha^{-1/2}(\sigma^y \otimes \mathbf{1}) - \alpha^{1/2}(\sigma^z \otimes \sigma^y) \\ \Delta(\alpha, \beta; T^Y) &= \beta^{-1/2}(\sigma^x \otimes \mathbf{1}) - \beta^{1/2}(\sigma^z \otimes \sigma^x) \\ \Delta(\alpha, \beta; E) &= \mathbf{1} \otimes \mathbf{1} + \mathbf{1} \otimes \mathbf{1}. \end{aligned} \quad (29)$$

It is trivial to check that similar to Eq. (18) we get

$$R \Delta(\alpha, \beta) R^{-1} = \Delta(\alpha^{-1}, \beta^{-1}). \quad (30)$$

Before concluding we would like to show that for $U_{\alpha, \beta}[SU(1/1)]$ one can introduce more than two parameters (as in the Lie algebra case when we had more than one). The most general chain which has a zero mode for all its values of the parameters is [5]

$$\begin{aligned} H_i &= \frac{1}{2} \{ \frac{\Theta + \Theta^{-1}}{2} (\eta \zeta \sigma_i^x \sigma_{i+1}^x + \eta^{-1} \zeta^{-1} \sigma_i^y \sigma_{i+1}^y) \\ &\quad + \frac{\Theta - \Theta^{-1}}{2} (\eta \zeta^{-1} \sigma_i^x \sigma_{i+1}^y + \eta^{-1} \zeta \sigma_i^y \sigma_{i+1}^x) \\ &\quad + q \sigma_i^z + q^{-1} \sigma_{i+1}^z \} \}. \end{aligned} \quad (31)$$

H_i depends on four parameters. One can check however that Eq. (21) holds with

$$\nu = \frac{q^2 + q^{-2}}{4} + \frac{(\eta^2 - \eta^{-2})(\zeta^2 - \zeta^{-2})}{8} + \frac{(\eta^2 + \eta^{-2})(\zeta^2 + \zeta^{-2})(\Theta^2 + \Theta^{-2})}{16}$$

$$\begin{aligned} \mu &= \left(\frac{(\eta^2 - \eta^{-2})(\zeta^2 - \zeta^{-2})}{4} + \frac{(\eta^2 + \eta^{-2})(\zeta^2 + \zeta^{-2})(\Theta^2 + \Theta^{-2})}{8} - 1 \right) \\ &\times \left(\frac{(q - q^{-1})^2}{2} + \frac{(\Theta - \Theta^{-1})^2 \left(\frac{\eta}{\zeta} - \frac{\zeta}{\eta} \right)^2}{8} \right) \end{aligned} \quad (32)$$

which means that we are back to two parameters.

To sum up we have shown that the two-parameter deformation of the $SU(1/1)$ superalgebra leads to a new Hopf algebra. One obtains then a generalisation of the Hecke algebra and of the braid group (see Eqs. (23) and (25) keeping the structure of connection between the R matrices and the coproduct (see Eq. (30)).

Acknowledgements: We would like to thank Franz Gaehlen for writing the computer program which made us discover Eq. (21). We would also like to thank D. Altschuler, L. Alvarez-Gaumé, D. Arnaudon, M. Chaichian, R. Coqueraux, C. Cinkovic and M. Scheunert for usefull discussions.

References

- [1] E. E. Demidov, Yu. I. Manin, E. E. Mukhin and D. V. Zhdanovich,
Progs. Theor. Phys. Suppl. **102** (1990) 203
A. Sudbery, *J. Phys. A: Math. Gen.* **23** (1990) L697
- [2] N. Yu. Reshetikhin, *Lett. Math. Phys.* **20** (1990) 331
- [3] N. den Nijs, in "Phase Transitions and Critical Phenomena",
Ed. C. Domb and J. L. Lebowitz, vol. 12, Academic Press 1988
- [4] E. D. Siggia, *Phys. Rev. B.* **16** (1977) 2319
- [5] H. Hinrichsen and V. Rittenberg, to be published
- [6] J. B. Kogut, *Rev. Mod. Phys.* **51** (1979) 659
- [7] M. Chaichian and P. Kulish, *Phys. Lett.* **B234** (1990) 72
E. Corrigan, D. Fairlie, P. Fletcher and R. Sasaki, *J. Math. Phys.* **31** (1990) 776
W. Schmidke, S. Vokos and B. Zumino, *Z. Phys.* **C48** (1990) 249
L. Kaufmann and H. Saleur, preprint EFI 90-42
L. Rozansky and H. Saleur, preprint YCTP-P20-91
- [8] H. Saleur, in "Trieste Conference on Recent Developements in Conformal Field Theories",
Eds. S. Randjekar-Daemi, E. Sezgin and J. B. Zuber, World Scientific 1989, p. 160
- [9] P. Martin and V. Rittenberg, preprint RIMS-770, 1991